

Coloured conical emission via second-harmonic generation

S. Trillo,^[†] C. Conti

Istituto Nazionale di Fisica della Materia (INFM)-RM3,
Via della Vasca Navale 84, 00146 Roma, Italy

P. Di Trapani, O. Jedrkiewicz, J. Trull ^[‡]

INFM and Dept. of Chemical, Physical and Mathematical
Sciences,
University of Insubria, Via Valleggio 11, 22100 Como, Italy

G. Valiulis

Dept. of Quantum Electronics, Vilnius University,
Building 3 Sauletekio Avenue 9, bldg. 3, LT-2040, Vilnius,
Lithuania

G. Bellanca

Dept. of Engineering, University of Ferrara, Via Saragat 1,
44100 Ferrara, Italy

Abstract

We predict that the combination of space and time modulational instabilities occurring via parametric wave-mixing in quadratic media leads to coloured conical emission. This phenomenon should be observed under conditions usually employed in second-harmonic generation experiments.

Modulational instability (MI) stands for the exponential growth of periodic, long-wavelength perturbations occurring either in time (i.e., temporal break-up [1]), or space (filamentation, e.g. Ref. [2]), due to interplay of Kerr nonlinearity and group-velocity dispersion (GVD) or diffraction, respectively. MIs occur also in quadratic media, in the form of parametric instabilities of two-color pump beams, encompassing both the cases of propagation invariant solutions (i.e., eigenmodes [3, 4]), and periodic pump evolutions [5, 6] as in, e.g. unseeded second-harmonic generation (SHG). Filamentation has been observed both in unseeded and seeded SHG geometries, in 1+2 (bulk) and 1+1 (planar waveguides) dimensions [4, 5, 7], while temporal MI have not been observed yet. On this ground, one might naturally wonder about the coupling of spatial and temporal degrees of freedom, as occurs, e.g., in self-focusing of short pulses. In this letter, our purpose is to show that, in the presence of both dispersion and diffraction, spatio-temporal MI in quadratic bulk samples cannot be simply seen as the mere superposition of filamentation and temporal break-up. Rather, it leads to a new phenomenon, namely coloured conical emission (CCE), where rings of different colours are expected to grow exponentially on top of both pump far-fields. As we show CCE via SHG has specific features which make it different from that earlier predicted for Kerr media [8, 9, 10]. It differs also from other space-time phenomena characteristic of SHG, such as snake instabilities [11], where the spatial breakup of SHG solitons associated with

temporal MI does not lead to substantial modifications of the beam angular spectra, which remain confined nearly on-axis.

Although the phenomenon of CCE implies component which travel off-axis, the angles usually involved are rather small and it is reasonable to use the following paraxial dimensionless model for SHG in noncritical configuration (i.e., no spatial walk-off)

$$\begin{aligned}
i \frac{\partial u_1}{\partial z} + \frac{\sigma_1}{2} \nabla_{\perp}^2 u_1 - \frac{\beta_1}{2} \frac{\partial^2 u_1}{\partial t^2} + u_2 u_1^* &= 0, \\
i \frac{\partial u_2}{\partial z} + \frac{\sigma_2}{2} \nabla_{\perp}^2 u_2 + i \delta \frac{\partial u_2}{\partial t} - \frac{\beta_2}{2} \frac{\partial^2 u_2}{\partial t^2} + \delta k u_2 + \frac{u_1^2}{2} &= 0,
\end{aligned} \tag{1}$$

Here, the (cw plane-wave) fields are conveniently normalized in terms of their total intensity I_t to yield $|u_2|^2 + |u_1|^2/2 = 1$. Accordingly the real-world distance $Z = zZ_{nl}$ is fixed by the nonlinear length scale $Z_{nl} = (\chi I_t)^{-1}$, where $\chi = k_0(2\eta_0/n_1^2 n_2)^{1/2} d_{eff}$ [3]. The normalized retarded time $t = (T - Z/V_{g1})/T_0$ and the Laplacian $\nabla_{\perp}^2 = \partial_x^2 + \partial_y^2$, where $(x, y) = (X, Y)/R_0$, are given in terms of temporal and transverse scales, $T_0 = (|k_1''|Z_{nl})^{1/2}$ and $R_0 = (Z_{nl}/k_1)^{1/2}$, respectively. Moreover $\sigma_m = k_1/k_m$, $m = 1, 2$ (we set $\sigma_2 = \frac{1}{2}$), $\beta_m = k_m''/|k_1''|$, $k_m'' = d^2k/d\omega^2|_{m\omega}$ are GVDs, $\delta k = (k_2 - 2k_1)Z_{nl}$ is the phase mismatch, and $\delta = Z_{nl}/Z_w$ stands for group-velocity mismatch (GVM), $Z_w = T_0(1/V_{g2} - 1/V_{g1})^{-1}$ being the walk-off length ($V_{gm} = dk/d\omega|_{m\omega}^{-1}$).

Let us consider propagation-invariant solutions of Eqs. (1), i.e. cw plane-wave eigenmodes $u_m = u_{m0}^{\pm} \exp(im\beta z)$, $m = 1, 2$, where $u_{20}^{\pm} = \beta = [\delta k \pm (\delta k^2 + 12)^{1/2}]/6$, $u_{10}^{\pm 2} = 2[1 - |u_{20}^{\pm}|^2]$, the upper and lower sign standing for in-phase ($\phi = \arg(u_2 - 2u_1) = 0$, existing for $-\infty < \delta k \leq 2$) and out-of-phase ($\phi = \pi$, $\infty > \delta k \geq -2$) modes, respectively [3]. We study the

stability of these waves by inserting in Eqs. (1) the fields

$$u_m(r, t, z) = [u_{m0} + a_m(r, t, z)] \exp(im\beta z), m = 1, 2, \quad (2)$$

where $u_{m0} = u_{m0}^\pm$, and we consider a radially symmetric perturbation:

$$a_m(r, t, z) = f_m^+(r, t) \exp(\lambda z) + f_m^-(r, t)^* \exp(\lambda^* z). \quad (3)$$

After linearization, we obtain a system of four partial differential equations in the unknowns $f_{1,2}^\pm(r, t)$. Without loss of generality, we introduce the Fourier-Bessel transform of $f_m^\pm$, $F_m^\pm(K, \Omega) = \mathcal{F}[f_m^\pm(r, t)]$, which permits to express the perturbation amplitudes through the inverse transform $f_m^\pm = \mathcal{F}^{-1}[F_m^\pm]$, as a superposition of conical J_0 Bessel waves [12] of transverse wavevector $K = [K_x^2 + K_y^2]^{1/2}$ and frequency Ω , as

$$f_m^\pm(r, t) = \int_{-\infty}^{\infty} \int_0^{\infty} F_m^\pm(K, \Omega) J_0(Kr) e^{i\Omega t} K dK d\Omega. \quad (4)$$

In terms of the vector $F = [F_1^+ \ F_1^- \ F_2^+ \ F_2^-]^T$ of transformed amplitudes, the linearized system becomes algebraic and takes the form

$$\begin{pmatrix} \Omega_1 + i\lambda & u_{20} & u_{10} & 0 \\ u_{20} & \Omega_1 - i\lambda & 0 & u_{10} \\ u_{10} & 0 & \Omega_2 - \delta\Omega + i\lambda & 0 \\ 0 & u_{10} & 0 & \Omega_2 + \delta\Omega - i\lambda \end{pmatrix} F = 0,$$

where $\Omega_m \equiv \beta_m \Omega^2 / 2 - \sigma_m K^2 / 2 - m\beta + (m-1)\delta k$. The compatibility condition for this 4×4 system gives the dispersion relation $\lambda = \lambda(K, \Omega)$. The eigenvalues λ have cumbersome expressions, except for vanishing GVM ($\delta = 0$ [13]), for which we find (set $2f_0 = u_{20}^2 - 2u_{10}^2 - \Omega_1 - \Omega_2$, $f_1 = (\Omega_1 \Omega_2 - u_{10}^2)^2 - u_{20}^2 \Omega_2^2$),

$$\lambda = \lambda(K, \Omega) = \pm \sqrt{f_0 \pm \sqrt{f_0^2 - f_1}}, \quad (5)$$

Instabilities sets in when at least one of the four eigenvalues λ in Eq. (5) have a positive real part, thus causing the perturbation to grow exponentially with gain $g = g(\Omega, K) = 2\text{Re}(\lambda)$. Two qualitatively different regimes are found depending on the sign of GVD. In Fig. 1 we display the typical spectral gain curve obtained in the anomalous GVD regime ($\beta_{1,2} = -1$). As shown, the gain is relatively narrowband, and its axisymmetric feature (g peaks at $\Omega^2 + K^2 = \text{constant}$) reflects the fact that space and time play indeed the same role. Conversely, when the GVD is normal, perturbations with large wavevector (angle) K and detuning Ω , can be amplified. Three different branches of growing perturbations exist even when the collinear process is phase-matched ($\delta k = 0$), as shown in Fig.2(a). This type of spatio-temporal MI entails CCE, which can be regarded as the amplification process of a superposition of Bessel J_0 beams characterized by progressively larger values of of angle-frequency ($K - \Omega$) pairs. The peak gain [obtained in a wide bandwidth of high frequencies $K - \Omega$, see Fig.2(a)], is displayed in Fig.2(b) and decreases monotonically for large $|\delta k|$ for both eigenmodes.

The underlying physics of CCE involve multiple three-photon processes. Pump photons at 2ω travelling on-axis decay into photon pairs at frequency $\omega - \delta\omega$ and $\omega + \delta\omega$ travelling off-axis with opposite angles. These in turn can produce off-axis photons at $2\omega \pm \delta\omega$ via sum-frequency mixing with on-axis pump photons at ω . In the limit $\delta k = \pm 2$ where the pump degenerate into the single component $u_{20}^\pm = \pm 1$ ($u_{10}^\pm = 0$), only the downconversion process takes place, and we can write the longitudinal projection (the transverse one is identically satisfied) of the noncollinear phase-matching condition as $k_z(2\omega) - k_z(\omega + \delta\omega) - k_z(\omega - \delta\omega) = 0$. In the paraxial approximation $k_z(\omega \pm \delta\omega) = [k(\omega \pm \delta\omega)^2 - k_t^2]^{1/2} \simeq k(\omega \pm \delta\omega) - k_t^2/2k_1$ (k_t is the real-world tranverse wavenumber) and by expanding $k(\omega \pm \delta\omega) = k_1 \pm V_{g1}^{-1}\delta\omega + k_1''\delta\omega^2/2$,

the phase-matching condition yields

$$k_1''\delta\omega^2 - k_1^{-1}k_t^2 = \Delta k. \quad (6)$$

In terms of dimensionless frequencies $K = k_t R_0$ and $\Omega = \delta\omega T_0$, Eq. (6) reads as $\Omega^2 - K^2 = \pm 2$, which turns out to be in perfect agreement with the locus $(K - \Omega)$ of peak gain as obtained from Eq. (5).

Remarkably we find that CCE is robust against the presence of GVM. In fact for $\delta \neq 0$ CCE occurs also in the ideal GVD-free limit ($\beta_{1,2} = 0$). The relative magnitude of GVM and GVD depends, however, on the intensity I_t . For instance, in a LBO crystal pumped at $\lambda_0 = 1 \mu\text{m}$ ($k_1'' = 0.045 \text{ ps}^2/\text{m}$, $k_2'' = 0.25 \text{ ps}^2/\text{m}$) with $I_t = 50 \text{ GW}/\text{cm}^2$ ($Z_{nl} = 0.6 \text{ mm}$), we find $Z_w = 0.12 \text{ mm}$ which yields $\delta = 5.4$. Figure 3 shows the different gain branches for this case, when the crystal operates in the large mismatch limit $k_2 - 2k_1 = -30 \text{ cm}^{-1}$ ($\delta k = -1.8$). The dimensional gain $G = g/Z_{nl}$ is reported against real-world angles $\theta \simeq \sin \theta = K/(k_1 R_0)$ and wavelength detunings $\Delta\lambda = \lambda_0^2 \Omega / (2\pi c T_0)$.

Though we have considered perturbations growing on top of the SHG eigenmodes, CCE can occur also from different launching conditions entailing exchange of energy between ω and 2ω pump beams. This case is similar to spatial or temporal MI occurring in unseeded SHG, and can be analyzed by means of Floquet techniques [5, 6]. In this letter, we rather present numerical evidence for such phenomenon, by integrating Eqs. (1) in 1+1+1D, setting $\partial_y = 0$. As an example Fig.4 shows the spectral content of the two fields after propagation ($z = 15$), as obtained when only the pump beam u_1 is launched at $z = 0$, on top of white noise. As shown, CCE favours the amplification of spatio-temporal frequencies which closely follow the spectral gain [see Fig.2(a)]. Due to the large bandwidth of the process we have

also found that CCE takes place from short pulse, narrow beam excitations, a case which will be discussed elsewhere.

We end up by emphasizing that CCE in SHG exhibits important differences with respect to CCE in true Kerr media. First, SHG makes CCE accessible in materials which are virtually GVD-free, and, in the large mismatch limit, for both signs of effective Kerr nonlinearity in the same physical material. However, the major importance of CCE stems from the fact that it can trigger the formation of novel spatio-temporal propagation-invariant wavepackets of Eqs. (1), i.e. $u_m(r, t, z) = q_m(r, t) \exp(im\beta z)$ [14]. In the normal GVD regime, $q_m(r, t)$ exist for $\beta < 0$ in the form of a polichromatic superposition of J_0 beams [12], so-called nonlinear X-waves [14], which exist also in the linear limit [15]. In SHG when the J_0 perturbations are amplified at the expense of an out-of-phase pump beam, they develop the right sign of nonlinear phase-shift ($\beta < 0$) for the formation of a nonlinear X-wave. Viceversa, in Kerr media, where nonlinear X-waves also exist, cw plane-waves and localized wavepackets experience opposite nonlinear phase-shifts, and such mechanism is unlikely to take place.

In summary we have shown that spatio-temporal MI of propagation-invariant as well as dynamically evolving pump beams in SHG gives rise to conical emission, which might play an important role in the space-time dynamics. Such phenomenon is intrinsic to quadratic materials and should not be confused with other recently reported mechanisms of conical emission [16].

Financial support from MIUR (PRIN project), INFN (PAIS project), UNESCO, MEC (Spain), and Lithuanian Science and Studies Foundation (grant T-491), is gratefully acknowledged. The work of C.C. is supported by the Fondazione Tronchetti Provera.

References

[†] Also with Dept. of Engineering, University of Ferrara.

[‡] Permanent Address: Dept. Física i Enginyeria Nuclear, UPC Terrassa, Spain

1. K. Tai, A. Hasegawa, and A. Tomita, Phys. Rev. Lett. **56**, 135 (1986).
2. R. Malendevich, L. Jankovic, G.I. Stegeman, and J. S. Aitchison, Opt. Lett. **26**, 1879 (2001).
3. S. Trillo and P. Ferro, Opt. Lett. **20**, 438 (1995).
4. R. Schiek, H. Fang, R. Malendevich, and G.I. Stegeman, Phys. Rev. Lett. **86**, 4528 (2001).
5. R.A. Fuerst, D.M. Baboiu, B. Lawrence, W.E. Torruellas, G.I. Stegeman, S. Trillo, and S. Wabnitz, Phys. Rev. Lett. **78**, 2756 (1997).
6. S. Trillo, and S. Wabnitz, Phys. Rev. E **55**, R4897 (1997).
7. H. Fang, R. Malendevich, R. Schiek, and G.I. Stegeman, Opt. Lett. **25**, 1786 (2000).
8. L.W. Liou, X.D. Cao, C.J. McKinstrie, and G.P. Agrawal, Phys. Rev. A **46** 4202 (1992).
9. G.G. Luther, A.C. Newell, J.V. Moloney, and E.M. Wright, Opt. Lett. **19** 789 (1994).
10. E.J. Fonseca, B. S. Cavalcanti, J.M. Hickmann, Opt. Commun. **169** 199 (1999).

11. A. De Rossi, S. Trillo, A.V. Buryak, and Yu.S. Kivshar, Phys. Rev. E **56**, R4959 (1997).
12. J. Durnin, J.J. Miceli, and J.H. Eberly, Phys. Rev. Lett. **58**, 1499 (1987).
13. R.J. Gehr, M.W. Kimmel, and A.V. Smith, Opt. Lett. **23**, 1298 (1998).
14. C. Conti *et al.* *Nonlinear X-waves: light bullets in normally dispersive media ?*, QELS Conference 2002, (Optical Society of America, Washington DC), paper QTuJ6.
15. P. Saari and K. Reivelt, Phys. Rev. Lett. **79**, 4135 (1997).
16. K.D. Moll, D. Homoelle, A.L. Gaeta, and R.W. Boyd, Phys. Rev. Lett. **88**, 153901 (2002).

Fig. 1. MI gain $g(\Omega, K)$ in the anomalous GVD regime for the in-phase eigenmode at phase-matching $\delta k = 0$.

Fig. 2. Normal GVD regime ($\beta_{1,2} = 1$): (a) Level plot of conical emission gain $g(\Omega, K)$ for $\phi = 0$ eigenmode (similar plot holds for $\phi = \pi$); (b) Peak value of g vs. δk .

Fig. 3. Level plot of CCE dimensional gain G for the in-phase eigenmode in a LBO crystal (see text for data).

Fig. 4. Spectrally selective amplification of white noise seen at $z = 15$ from unseeded collinearly phase-matched SHG [$\delta k = 0$ in Eqs. (1)] pumped by a cw plane-wave pump at ω : (a) $|u_1(K, \Omega)|$; (b) $|u_2(K, \Omega)|$.

This figure "fig1OL_CCE.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/physics/0204042v2>

This figure "fig2OL_CCE.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/physics/0204042v2>

This figure "fig3OL_CCE.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/physics/0204042v2>

This figure "fig4OL_CCE.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/physics/0204042v2>